## Molecular Geometry

How can molecular shapes be predicted using the VSEPR theory?

### Why?

When you draw a Lewis structure for a molecule on paper, you are making a two-dimensional representation of the atoms. In reality however, molecules are not flat—they are *three*-dimensional. The true shape of a molecule is important because it determines many physical and chemical properties for the substance. In this activity you will learn how to predict molecular shapes.

### Model 1 – Lewis Structures



1. Name the type of structures shown in the left-hand column of Model 1.

### Lewis Structures

- 2. Examine the drawings in Model 1.
  - a. What does a solid line between two element symbols represent in the drawings of the molecules? Bond
  - b. What subatomic particles (protons, neutrons or electrons) make up these solid lines? electrons
  - c. What does a pair of dots represent in the drawing of the molecules?



- d. What subatomic particle (protons, neutrons or electrons) makes up each dot? electrons
- 3. In the English language, what does the word "domain" mean? (Your group must come to consensus on this question.)
- 4. Which molecules in Model 1 have four electron domains? Circle or highlight the four electron domains in the Lewis structure for each molecule that you identified.
  3.4.5
- 5. Which molecules in Model 1 have two electron domains? Circle or highlight the two electron domains in the Lewis structure for each molecule that you identified.

3045 2,6

- Which molecule in Model 1 has three electron domains? Circle or highlight the three electron domains in the Lewis structure for the molecule that you identified.
- 7. When determining the number of electron domains in a Lewis structure, which of the following should you count? Find evidence from Model 1 to support your answers.

*a.* Bonds on the center atom

(b) Lone pairs on the center atom

c. Total number of atoms in the molecule indirectly because it affects the number of bonds

KLone pairs on peripheral atoms

8. When determining the number of electron domains in a Lewis structure, do you count double bonds as one domain or two domains? Find evidence to support your answer from Model 1.

one domain #6 only has 2 domains and it has 2 double bonds

9. Explain the difference between a **bonding electron domain** and a **nonbonding electron domain** using the examples in Model 1.

```
bonding electron domain - bond - shared pair of
electrons
```

non bonding domain - lone pair - un shaved electrons 10. Circle the correct word or phrase to complete the sentences:

Pairs of electrons will (attract/tepe) each other.

Two bonds on the same atom will try to get as (close to/far from) each other as possible.

A lone pair of electrons and a bonded pair of electrons will (push away from/move toward) each other.

# Read This!

STO

The VSEPR (Valence Shell Electron Pair Repulsion) Theory helps predict the shapes of molecules and is based on the premise that electrons around a central atom repel each other. Electron domains are areas of high electron density such as bonds (single, double or triple) and lone-pairs of electrons. In simple terms VSEPR means that all electron bonding domains and electron nonbonding domains around a central atom need to be positioned as far apart as possible in *three-dimensional* space.

11. VSEPR theory specifies "valence shell" electrons. Explain why these are the most critical electrons for determining molecular shape based on your exploration of Model 1.

these are the electrons involved in bonding

- 12. In the VSEPR theory, what is repelling what? electron pairs
- 13. Based on the information in the *Read This!* section, sketch one of the molecular shapes shown below in each of the boxes provided in Model 1.

	Linear	Trigonal planar	
Three-Dimensional Molecular Shapes	180°		
Tetrahedral	Pyramidal Pyramidal	Bent 104.5°	



14. Often we draw Lewis structures with 90° bond angles. Do any of the molecular shapes in Model 1 have 90° bond angles?

no

15. Are the bond angles in the three-dimensional molecules generally larger or smaller than those shown in the Lewis structures drawn on notebook paper?

16. Why is it possible to get larger angles separating electron domains in three-dimensions versus two-dimensions?

you can use the space infront/behind the paper

- 17. Identify the three molecules shown in Model 1 that have four electron domains each.
  - a. What happens to the size of the bond angle(s) in a molecule as the number of lone pairs on the central atom increases?

b. Discuss in your group some possible explanations for the trend in part a. Your presenter should be ready to present to the class one or two of your hypotheses for full class discussion.

## STOP

- 18. A student does not "waste" his time drawing a Lewis structure before determining the shape of PF<sub>3</sub>. The student thinks that the shape of PF<sub>3</sub> must be trigonal planar because there are three fluorine atoms bonded to the central phosphorus atom.
  - a. Draw the Lewis structure for  $PF_3$ .

5+3(7)-26

b. Was the student's answer for the shape of a PF, molecule correct? Explain.

### No

c. Why is it important to draw the Lewis structure for a molecule before identifying the shape of the molecule?

To be able to count the bonding and nonbonding

19. Complete the following chart:

Molecule	Lewis Structure	3-D Drawing	Name of 3-D Shape	Bond Angle
H <sub>2</sub> S	H-S-H	0 0	bent	104.5
PH <sub>3</sub>	H - P - H	0.0	trigonal pyramidal	107.
$\operatorname{CCl}_4$	: ci : : ci : : ci :	0	tetrahedral	109.5
CS <sub>2</sub>	Ŝ=C=Ŝ	0=0=0	linear	180.

STOP

#### **Extension Question**

- 20. Ozone, O3, is not a linear molecule. Actually it is bent with an angle that is a little less than 120°.
  - a. Draw the Lewis structure of ozone, O3.

b. Describe why ozone has a bent shape instead of a linear shape.

c. Describe why ozone's bond angle is larger than that of water, H<sub>2</sub>O.

¢